

Photon statistics of high- β quantum dot photonic-crystal lasers

Y.-S. Choi¹, M. T. Rakher², S. Strauf^{2,3}, K. Hennessy¹, A. Badolato¹,
D. Bouwmeester², P. M. Petroff^{1,3}, and E. L. Hu^{1,3}

¹ECE, ²Physics, ³Materials, University of California, Santa Barbara, California 93106

We demonstrate the photon statistics of InAs quantum-dot (QD) photonic-crystal (PC) lasers with high spontaneous emission coupling factors, $\beta > 0.25$. Such high β factors lead to the onset of lasing at incident pump powers $< 40 \text{ W/cm}^2$ and pronounced linewidth narrowing above threshold. Measurements of the second-order correlation function, $g^{(2)}(T)$, confirm that a transition of the photon statistics from random spontaneous emission to coherent stimulated emission occurs. We then use the Fano-Mandel parameter^[1], $F = [\langle a^\dagger a a^\dagger a \rangle - \langle a^\dagger a \rangle^2] / \langle a^\dagger a \rangle$, to demonstrate the first direct identification of lasing threshold in high- β , soft-turn-on devices.

With further optimization of QD placement using active positioning schemes^[2], we expect to realize a thresholdless laser operating on a single QD.

[1] R. Loudon, *The Quantum Theory of Light* (Oxford University, New York, 2000)

[2] A. Badolato, K. Hennessy, M. Atature et. al. in publication, *Science* (2005).